



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

STANDARD BOOKS

Plane and Spherical Trigonometry

Revised and Enlarged Edition

By GEORGE N. BAUER and W. E. BROOKE, University of Minnesota.

THE new edition contains more problems and embodies such modifications as have been suggested by experience in the classroom. There have also been added

Logarithmic and Trigonometric Tables

Including a few three-place, four-place, and complete five-place tables. The tables fill 140 pages. Price of Trigonometry with Tables, \$1.60. Tables separately, 64 cents.

Analytic Geometry

By W. A. WILSON and J. I. TRACEY, Department of Mathematics, Yale University.

THIS book presents in a short course those parts of Analytic Geometry which are essential for the study of Calculus. The material has been so arranged that topics which are less important may be omitted without a loss of continuity. The text is therefore adapted for use in classes which aim to cover in one year the fundamental principles and applications of both Analytic Geometry and Calculus. Cloth. x+212 pages. Price, \$1.28.

Fite's College Algebra

THE clearness, brevity, and rigor of this book won for it widely extended use from the day of its publication. Its perfect adaptation to the needs of college classes is indicated by its steadily increasing sale. 289 pages. \$1.48.

Miller and Lilly's Analytic Mechanics

A course that is distinctly teachable, practical, rigorous, and adaptable. Abundant problems and exercises are included. 312 pages. \$2.00.

Correspondence invited

D. C. HEATH & COMPANY, Publishers

Boston

New York

Chicago

London

School Science and Mathematics

A Monthly Journal for all Science and Mathematics Teachers

It is especially Interesting and Helpful to all Mathematics Teachers in Secondary Schools and to all other Instructors in Mathematics who wish to keep in close touch with the latest Thought and Ideas in High School Mathematics.

Mathematics Department Edited by Professor Herbert E. Cobb, Head of Mathematics Department, Lewis Institute, Chicago. Problem Department Edited by Dr. J. O. Hassler, Crane Junior College and High School, Chicago.

Subscribe now

\$2.50 per year

School Science and Mathematics

2059 East 72nd Place

CHICAGO

Ready about January 1, 1918

ANALYTIC GEOMETRY

By EDWIN S. CRAWLEY and HENRY B. EVANS

Professors of Mathematics in the University of Pennsylvania

Size: xiv+239 pages, $7\frac{1}{4} \times 4\frac{3}{4}$ inches. Price \$1.60.

Chapters I to X (190 pages) give a full college course in plane analytic geometry. Chapter XI (14 pages) on empirical equations will be of particular interest to students of engineering and other applied sciences. Chapter XII, the concluding chapter, is devoted to the extension of coordinate geometry to some space problems.

Orders and applications for sample copies for examination with a view to introduction should be addressed to

E. S. CRAWLEY, University of Pennsylvania, Philadelphia

Teachers of Mathematics

SHOULD READ

The Mathematics Teacher

The only journal in America devoted entirely to the interests of the teaching of mathematics. It is helping hundreds of others and will help you.

No teacher of mathematics should be without it and you will not be, if a progressive teacher.

Subscription Price, \$1.00 a year

THE MATHEMATICS TEACHER

103 Avondale Place

SYRACUSE, NEW YORK

THE NEW ERA PRINTING COMPANY

LANCASTER, PA.

Is prepared to execute in first-class and satisfactory manner all kinds of printing and electrotyping. Particular attention given to the work of Schools, Colleges, Universities, and Public Institutions.

Books, Periodicals

Technical and Scientific Publications

Monographs, Theses, Catalogues

Announcements, Reports, etc.

All Kinds of Commercial Work

(Printers of the Bulletin and Transactions of the American Mathematical Society, etc., etc.)

Publishers will find our product ranking with the best in workmanship and material, at satisfactory prices. Our imprint may be found on a number of high-class Technical and Scientific Books and Periodicals. Correspondence solicited. Estimates furnished.

THE NEW ERA PRINTING COMPANY

THE MULTIPLEX WRITING MACHINE

Special Mathematical Model

Two complete sets on one machine,—of any Science or Language, or many correspondence faces. All printable on one machine. "Many typewriters in one." "Just Turn the Knob."

Sample Problem

To solve $\frac{\partial^2 \varphi}{\partial t^2} = \sqrt{1+(\Delta h)^2} \frac{\partial^2 \varphi}{\partial x^2}$, put $m^2 = \sqrt{1+(\Delta h)^2}$ and assume $\varphi = \tau(t) \cdot \xi(x)$; so that

$$\frac{\partial^2 \varphi}{\partial t^2} = \frac{d^2 \tau}{dt^2} \cdot \xi(x), \text{ and } \frac{\partial^2 \varphi}{\partial x^2} = \tau(t) \frac{d^2 \xi}{dx^2}. \quad \text{Sub-}$$

stituting these into the original equation, we find that the variables, t and x , can be separated by dividing through by $\tau \cdot \xi$ where-

$$\text{upon we have } \frac{d^2 \tau}{dt^2} \div \tau = m^2 \frac{d^2 \xi}{dx^2} \div \xi. \quad \text{Since the}$$

first of these two equal members cannot vary when t changes, nor the second when x changes, both must remain equal to some constant, say,

$-m^2 n^2$. The two resulting equations yield the solutions

$$\xi = K_1 \cdot \sin[nx + \beta_1], \quad \tau = K_2 \cdot \sin[mnt + \beta_2]$$

$$\text{whence } \varphi = K_1 K_2 \sin[nx + \beta_1] \sin[mnt + \beta_2]$$

which we may then reduce to a more useful form:

$$\varphi = \sum_{n=0}^{n=\infty} A_n \sin[n(x \pm mt) + \delta_n].$$

An interesting fallacy results from applying the method of integration by parts, $\int u \cdot dv = uv - \int v \cdot du$, to a case where $u = 1/x$ and $dv = dx$: we get

$$\begin{aligned} \int \frac{dx}{x} &= \frac{1}{x} \cdot \int dx - \int x \cdot \{-1/x^2\} \\ &= 1 + \int dx/x \quad \text{whence } 0=1 !! \end{aligned}$$

$$\begin{aligned} \int \frac{dx}{5+7x^2} &= 1/5 \int \frac{dx}{1+\frac{7}{5}x^2} = \frac{1}{5} \sqrt{\frac{5}{7}} \int \frac{\sqrt{\frac{7}{5}} dx}{1+[\frac{7}{5}x]^2} \\ &= \frac{1}{35} \arctan [\sqrt{7/5} x]. \end{aligned}$$

THE HAMMOND TYPEWRITER COMPANY

Actual Facsimile by Prof. Ransom of Tufts College. Special Booklet on Application.

604 East 69th St., New York City, N. Y.